

Severe Yeast Infection Protocol

Severe Yeast Infections

The following is for severe cases and can be altered by the treating physician. All simple sugars should be stopped, as refined carbohydrates feed yeast. See attached anti-inflammatory diet.

WEEKS 0-2

ADP 3-5 tid, 3 for a smaller person, 5 for someone over 250 pounds, (3 times a day with or without food) - kills fungus and a host of bacteria that support yeast growth.

Caprin 3 tid – coats the colon in a way that restricts the colonization and growth of yeast.

Beta TCP 3 tid – this will thin and mobilize your bile, which changes the environment of the bowel to discourage dysbiotic flora (yeast).

TolerAid 1 bid (2 times a day) breaks down the chemicals that the yeast put off, which make us feel sick.

Adult ENT Pro 1 bid – this formula has been shown to adhere to the mucosal linings of the mouth and sinus cavity and kills 3 different types of yeast. It also enhances the immune system to fight all forms of bacteria and yeast.

WEEKS 2-4

Changing dysbiotic agents is common in medicine. Changing botanicals will often kill resistant strains that were able to withstand the environmental stress that the earlier botanicals.

FC Cidal 2 tid – kills fungal forms that hide deep in tissue.

Dysbiocide 2 tid – designed for small worms. It has antibacterial and anti-fungal properties.

Beta TCP 3 tid

TolerAid 1 bid

BioDop7 Plus 1 at breakfast, 2 at bed to reestablish healthy bacterial flora.

WEEKS 5-6

ADP 2 tid

Berberine HCL 2 tid

Beta TCP 3 tid

TolerAid 1 bid (2 times a day) breaks down the chemicals that the yeast put off which make us feel sick.

Adult ENT Pro 1 bid

Severe Yeast Infection Protocol

WEEKS 7-8

FC Cidal 2 tid kills fungal forms that hide deep in tissue.

Dysbiocide 2 tid allow designed for small worms it has antibacterial and anti-fungal properties.

Beta TCP 3 tid

TolerAid 1 bid

BioDop7 Plus 1 breakfast 2 at bed to reestablish healthy bacterial flora.

VAGINAL DOUCHE

If the yeast or fungal infection is in the genital urinary tract, add the following as a vaginal douche: two to three ounces of distilled water, add Bio-AE Mulsion 10 drops, Liquid Iodine Forte 100 drops, and BioDop7 Plus 1 capsule. Use the douche every other day for 14 days. Use a rectal syringe for the douche and try to retain it for 7 to 8 minutes.

If extremely chronic, consider buying new underwear or at least dampen and put in a microwave to kill any yeast that may be present.

If recurrent, treat sexual partner that is reintroducing infection.